
LATEX MICRO

Sterile Nitrile Coating



Boost tactile sensitivity

- **Stretches and conforms** to your hand contour, staying comfortably in place
- Ideal in a thin double-gloving system where fingertip sensation is essential
- 30 percent thinner for enhanced flexibility and **tactile sensitivity**¹
- Interlocking, beaded cuff design helps to prevent roll-down
- Proprietary hand mold with an independent thumb design allows for an anatomical fit and natural movement



Backed by the expertise of Cardinal Health, Protexis™ Latex Micro Surgical Gloves meet all relevant FDA and ASTM standards, including also the requirements of European harmonized standards EN 455-1, -2, -3 and-4. They include those for physical dimensions, physical properties and freedom from holes.

Documentation is available upon request.

Designed for procedures where tactile sensitivity is the highest priority, Protexis™ Latex Micro Surgical Gloves provide a 'bare hand' feel, excellent tactile response and a smooth, anti-slip finish.



PRODUCT INFORMATION | LATEX MICRO

Properties (before aging) ⁷		
Force at break	17N³	
elongation at break (elasticity)	≥ 750%4	
Puncture resistance (cuff)	Min 5N⁵	
Freedom from holes	0.65 AQL ⁶	

When chemotherapy drugs are present, glove selection should be based on the specific type(s) of chemicals used. Users must refer to the instructions for use supplied with the box, for the chemicals being used to determine an adequate level of protection.

Chemotherapy agent permeation ¹⁰		
	Agent	Minimum breakthrough detection time in minutes (0.01 µg/cm²/minute)
1	Carmustine (3.3 mg/mL)	12.5 (Do not use)
2	Cisplatin (1.0 mg/mL)	> 240
3	Cyclophosphamide (20 mg/mL)	> 240
4	Doxorubicin HCL (2.0 mg/mL)	> 240
5	Etoposide (20 mg/mL)	> 240
6	Fluorouracil (50 mg/mL)	> 240
7	Ifosfamide (50 mg/mL)	> 240
8	Methotrexate (25 mg/mL)	> 240
9	Mitomycin C (0.5 mg/mL)	> 240
10	Mitoxantrone (2.0 mg/mL)	> 240
11	Paclitaxel (6.0 mg/mL)	> 240
12	Thiotepa (10 mg/mL)	22.9 (Do not use)
13	Vincristine Sulfate (1.0 mg/mL)	> 240





- 1. As compared to Protexis(TM) Latex Surgical Gloves
- 2. Caution: This product contains natural rubber latex which may cause allergic reactions, including anaphylactic response
- 3. Force at Break tested in accordance with EN 455-2
- 4. Elongation at Break tested in accordance with ASTM D 3577
- 5. Puncture resistance (cuff) tested in accordance with AS/NZS 4179
- 6. Freedom from holes tested in accordance with EN 455-1
- 7. Thickness tested in accordance with ASTM D 3577
- 8. Protein content tested in accordance with EN 455-3 using ASTM D 5712
- 35% reduction of materials used as compared to previous Cardinal Health packaging design. Cardinal Health Technical Council Conference 2011, presentation on file
- 10. Chemotherapy agent permeation tested in accordance with ASTM D 6978-05

THE CARDINAL HEALTH™ PROTEXIS™ SURGICAL GLOVES PROMISE:

We protect so you can perform.

Cardinal Health is dedicated to providing protection, performance and expertise so wearers can perform confidently and focus on their patients.



Help maximize storage space: Half-fold packaging design reduces packaging material9.



Storage recommendations: Do not store near sources of ultraviolet light, heat or x-rays. Store in a cool, dry place, store away from direct sunlight.



Expiration: 35 months from date of manufacture. Expiration date is printed on packaging.

For healthcare professionals only

Important information: Prior to use, refer to the instructions on the dispenser box supplied with this device for indications, contraindications, side effects, suggested procedure, warnings and precautions. As part of its continuous product development policy, Cardinal Health reserves the right to change product specifications without prior notification Please contact your Cardinal Health representative for additional product availability information.



© 2018 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE and PROTEXIS are trademarks of Cardinal Health and may be registered in the US and/or in other countries. Lit. No. 2CHL18-704524 (10/2018)